Salsola laricina

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **6**
Moisture indicator value: **5.4**
Reaction indicator value: **7.9**
Nutrient indicator value: **5**
Salinity indicator value: **5.8**

Disturbance indicator values

Disturbance frequency: **2.46**
Disturbance frequency (herb layer): **2.39**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.16**
Grazing pressure: **0.49**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **EC (KAL) *Kalidietea foliati***

Broad habitat

Occurrence in broad habitats: **Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz