

# *Salvia fruticosa*

## Habitus and growth type

Plant height [m]: **0.85**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]:

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **9.5**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **2.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **0.24**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.06**

Grazing pressure: **0.3**

Soil disturbance: **0.17**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) \*Ononido-Rosmarinetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#)

Constant species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**