

# *Salvia viridis*

## Habitus and growth type

Plant height [m]: **0.34**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **26.54**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **2.44**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9.5**

Moisture indicator value: **2.5**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.13**

Disturbance frequency (herb layer): **2.03**

Disturbance severity: **0.79**

Disturbance severity (herb layer): **0.62**

Mowing frequency: **0.28**

Grazing pressure: **0.33**

Soil disturbance: **0.35**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S72 Eastern Mediterranean spiny heath \(\*phrygana\*\)](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Synanthropic**