

Saponaria pumila

Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.02**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **1.7**

Moisture indicator value: **3.5**

Reaction indicator value: **1.5**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.44**

Disturbance frequency (herb layer): **1.44**

Disturbance severity: **0.3**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0**
Grazing pressure: **0.23**
Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CT \(TRI\) *Juncetea trifidi*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),
Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **1**