

Sarcocornia fruticosa

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **August-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9**

Moisture indicator value: **8**

Reaction indicator value: **8.5**

Nutrient indicator value: **7**

Salinity indicator value: **8**

Disturbance indicator values

Disturbance frequency: **1.85**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.01**

Grazing pressure: **0.21**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MG \(SAL\) *Salicornietea fruticosae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA253 Mediterranean mid-low saltmarsh](#)

Constant species of EUNIS habitats: [MA251 Mediterranean upper saltmarsh](#), [MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA253 Mediterranean mid-low saltmarsh](#)

Dominant species of EUNIS habitats: [MA253 Mediterranean mid-low saltmarsh](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Saline vegetation**