

# *Satureja obovata*

## Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **8.75**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8.8**

Moisture indicator value: **2.8**

Reaction indicator value: **6.7**

Nutrient indicator value: **3.1**

Salinity indicator value: **0.6**

### Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.07**

Grazing pressure: **0.39**

Soil disturbance: **0.15**

## Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Scrub**