Saussurea discolor

Habitus and growth type

Plant height [m]: **0.13**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.9**
Temperature indicator value: **1.6**
Moisture indicator value: **5.5**
Reaction indicator value: **7.7**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.07**
Disturbance frequency (herb layer): **1.28**
Disturbance severity: **0.3**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.19**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz