# *Saxifraga conifera*

## Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.03**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **3**
Moisture indicator value: **2**
Reaction indicator value: **7**
Nutrient indicator value: **1**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.86**
Disturbance frequency (herb layer): **2.2**
Disturbance severity: **0.41**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.21**
Grazing pressure: **0.41**
Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **FP (ONO) *Festuco hystricis-Ononidetea striatae***

### EUNIS habitat

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Diagnostic species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**

Constant species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*