Saxifraga fragosoi

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **5**
Moisture indicator value: **2**
Reaction indicator value: **7**
Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.88**
Disturbance frequency (herb layer): **0.94**
Disturbance severity: **0.3**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.14**
Grazing pressure: **0.2**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea***, **KC (ASP) *Asplenietea trichomanis***, **KG (PHA) *Phagnalo saxatilis-Rumicetea indurati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U33 Temperate, lowland to montane siliceous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

inland cliff, **U34 Mediterranean siliceous inland cliff**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz