Scabiosa taygetea

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-May**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **8**

Disturbance indicator values

Disturbance frequency: **0.89**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.09**
Grazing pressure: **0.32**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

vegetated (incl. rock and scree)

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz