

Scleranthus polycarpus

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **44.56**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **1.71**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**

Moisture indicator value: **3**

Nutrient indicator value: **1.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.06**

Disturbance frequency (herb layer): **2.18**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.49**

Grazing pressure: **0.37**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*, FI \(TUB\) *Helianthemetea guttati*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **3**