

# *Scorzoneroides carpetana*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **4**

Moisture indicator value: **4**

Reaction indicator value: **3**

Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **2.17**

Disturbance frequency (herb layer): **2.17**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.7**

Grazing pressure: **0.55**

Soil disturbance: **0.3**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1N Open Iberian supramediterranean dry acid and neutral grassland](#), [R33 Mediterranean short moist grassland of mountains](#)

Constant species of EUNIS habitats: [R33 Mediterranean short moist grassland of mountains](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**