

Scrophularia lucida

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8.7**

Moisture indicator value: **2.5**

Reaction indicator value: **7.3**

Nutrient indicator value: **3.3**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.64**

Mowing frequency: **0.36**

Grazing pressure: **0.26**

Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U28 Western Mediterranean base-rich scree](#),
[U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**