Scrophularia peregrina

Habitus and growth type

Plant height [m]: **0.56**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.7**
Temperature indicator value: **9**
Moisture indicator value: **5**
Reaction indicator value: **6.7**
Nutrient indicator value: **6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.67**
Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.81**
Disturbance severity (herb layer): **0.68**
Mowing frequency: **0.5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.21**
Soil disturbance: **0.38**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea*, QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz