

# *Scutellaria galericulata*

## Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **39.55**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.72**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **5.3**

Moisture indicator value: **9**

Reaction indicator value: **6.6**

Nutrient indicator value: **5.9**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.28**

Grazing pressure: **0.15**

Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [IA \(ALN\) \*Alnetea glutinosae\*](#), [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T15 Broadleaved swamp forest on non-acid peat](#)

Constant species of EUNIS habitats: [S92 \*Salix fen scrub\*](#), [T12 \*Alnus glutinosa\*-\*Alnus incana\* forest on riparian and mineral soils](#), [T15 Broadleaved swamp forest on non-acid peat](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality amplitude: **8**