Sedum acre

Habitus and growth type

Plant height [m]: **0.09**
Life span: **Perennial**
Life form: **Chamaephyte**

Leaf

Specific leaf area [mm^2/mg]: **10.9**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**
Moisture indicator value: **2.1**
Reaction indicator value: **6.6**
Nutrient indicator value: **1.7**
Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.29**
Disturbance frequency (herb layer): **1.77**
Disturbance severity: **0.46**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.33**
Grazing pressure: **0.31**
Soil disturbance: **0.33**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, CK (SED) *Sedo-Scleranthetea*, DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N15 Atlantic and Baltic coastal dune grassland (grey dune), R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops, R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops, R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**

Constant species of EUNIS habitats: **N15 Atlantic and Baltic coastal dune grassland (grey dune), R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops, R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops, R16 Perennial rocky grassland of Central and South-Eastern Europe, R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz