Sedum rubens

Habitus and growth type

Plant height [m]: **0.08**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **32.86**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **7.4**
Moisture indicator value: **2.8**
Reaction indicator value: **5**
Nutrient indicator value: **2.8**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.91**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.31**
Grazing pressure: **0.34**
Soil disturbance: **0.34**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, FI (TUB) *Helianthemetea guttati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz