# *Senecio gallicus*

## Habitus and growth type

Plant height [m]: **0.25**
Life span: **Annual**
Life form: **Therophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **19.27**

## Flower

Flowering period: **February-June**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **7.7**
Moisture indicator value: **2**
Reaction indicator value: **5**
Nutrient indicator value: **3.5**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.11**
Disturbance frequency (herb layer): **1.85**
Disturbance severity: **0.6**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity (herb layer): **0.54**
Mowing frequency: **0.35**
Grazing pressure: **0.28**
Soil disturbance: **0.37**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*