

Senecio sylvaticus

Habitus and growth type

Plant height [m]: **0.38**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **29.5**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.25**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **5.5**

Reaction indicator value: **4.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.51**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.16**

Grazing pressure: **0.22**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) Epilobietea angustifolii](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#)

Constant species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#),
[R1L Madeiran oromediterranean siliceous dry grassland](#), [R57 Herbaceous forest clearing vegetation](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

Distribution

Continentality: **4**

Continentality amplitude: **4**