Senecio transiens

Habitus and growth type

Plant height [m]: **0.29**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **January-May, October-December**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **8**
Moisture indicator value: **5**
Reaction indicator value: **7**
Nutrient indicator value: **7**
Salinity indicator value: **5**

Disturbance indicator values

Disturbance frequency: **1.41**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.14**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.2**
Soil disturbance: **0.52**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **N22 Mediterranean and Black Sea coastal shingle beach**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz