

Sesamoides interrupta

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **5**

Reaction indicator value: **2**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **2.1**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.09**

Grazing pressure: **0.34**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FM \(PIL\) *Saginetea piliferae***](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**