

Sesleria caerulea

Habitus and growth type

Plant height [m]: **0.27**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **19.27**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.12**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **3.2**

Moisture indicator value: **4.4**

Reaction indicator value: **8.3**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.47**

Disturbance frequency (herb layer): **1.22**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.12**

Grazing pressure: **0.24**

Soil disturbance: **0.11**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R44 Arctic-alpine calcareous grassland](#), [T36](#)

[Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Constant species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [R16 Perennial rocky](#)

[grassland of Central and South-Eastern Europe](#), [R18 Perennial rocky calcareous](#)

[grassland of subatlantic-submediterranean Europe](#), [R1A Semi-dry perennial](#)

[calcareous grassland \(meadow steppe\)](#), [R44 Arctic-alpine calcareous grassland](#),

[R51 Thermophilous forest fringe of base-rich soils](#), [S21 Subarctic and alpine](#)

[dwarf Salix scrub](#), [S22 Alpine and subalpine ericoid heath](#), [S24 Subalpine](#)

[genistoid scrub of the Amphi-Adriatic region](#), [S26 Subalpine Pinus mugo scrub](#),

[T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#), [T36](#)

[Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#),

[U36 Temperate high-mountain base-rich inland cliff](#), [U37 Temperate, lowland to](#)

[montane base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: [Coastal cliff](#), [Mire](#), [Grassland \(non-alpine, non-](#)

[saline\)](#), [Alpine-subalpine and arctic grassland](#), [Heathland](#), [Oromediterranean](#)

[scrub](#), [Scrub](#), [Forest](#), [Sparsely vegetated \(incl. rock and scree\)](#)

Distribution

Continentality: **3**

Continentality amplitude: **3**