

# *Sesleria tenerrima*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **7.1**

Moisture indicator value: **4.1**

Reaction indicator value: **7**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.05**

Grazing pressure: **0.18**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#), [FR \(DAP\) \*Daphno-Festucetea\*](#), [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1K \*Balkan and Anatolian oromediterranean dry grassland\*](#)

Constant species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Dominant species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**