

# *Sesleria vaginalis*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7.9**

Moisture indicator value: **3.5**

Reaction indicator value: **7**

Nutrient indicator value: **3.4**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.91**

Disturbance frequency (herb layer): **2.05**

Disturbance severity: **0.39**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.02**

Grazing pressure: **0.33**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[FR \(DAP\) Daphno-Festucetea](#)**

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**