

Schedonorus pratensis

Habitus and growth type

Plant height [m]: **0.64**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **27.6**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.02**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **4.5**

Reaction indicator value: **5.9**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.77**

Disturbance frequency (herb layer): **2.11**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **1.3**

Grazing pressure: **0.27**

Soil disturbance: **0.33**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R21 Mesic permanent pasture of lowlands and mountains](#), [R22 Low and medium altitude hay meadow](#), [R34 Submediterranean moist meadow](#)

Constant species of EUNIS habitats: [R21 Mesic permanent pasture of lowlands and mountains](#), [R22 Low and medium altitude hay meadow](#), [R23 Mountain hay meadow](#), [R34 Submediterranean moist meadow](#), [R35 Moist or wet mesotrophic to eutrophic hay meadow](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**, **Alpine-subalpine and arctic grassland**, **Saline vegetation**, **Scrub**