Sideritis taurica

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.1**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.3**
Moisture indicator value: **5.3**

Disturbance indicator values

Disturbance frequency: **2.09**
Disturbance frequency (herb layer): **2.09**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.51**
Mowing frequency: **1.11**
Grazing pressure: **0.36**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KH (DRY) *Drypidetea spinosae***

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz