

# *Sisymbrium irio*

## Habitus and growth type

Plant height [m]: **0.34**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **20.14**

## Flower

Flowering period: **January-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3.8**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.85**

Disturbance frequency (herb layer): **2**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.87**

Grazing pressure: **0.19**

Soil disturbance: **0.57**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) \*Sisymbrietea\*](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

## **Distribution**

Continentality: **4**

Continentality amplitude: **4**