Sorbus aucuparia

Habitus and growth type

Plant height [m]: **9.27**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **14.45**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **205.69**
Dispersal mode: **Endozoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.4**
Moisture indicator value: **5.8**
Nutrient indicator value: **3.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.07**
Disturbance frequency (herb layer): **1.25**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.15**
Mowing frequency: **0.01**
Grazing pressure: **0.19**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1F Baltic coniferous coastal dune forest**, **T18 Fagus forest on acid soils**, **T31 Temperate mountain Picea forest**, **T32 Temperate mountain Abies forest**, **T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest**, **T35 Temperate continental Pinus sylvestris forest**, **T3F Dark taiga**, **T3G Pinus sylvestris light taiga**, **T3H Larix light taiga**, **T3K Picea mire forest**

Constant species of EUNIS habitats: **N1D Atlantic and Baltic broad-leaved coastal dune forest**, **N1F Baltic coniferous coastal dune forest**, **R52 Forest fringe of acidic nutrient-poor soils**, **R57 Herbaceous forest clearing vegetation**, **S25 Subalpine and subarctic deciduous scrub**, **S26 Subalpine Pinus mugo scrub**, **S32 Temperate Rubus scrub**, **S38 Temperate forest clearing scrub**, **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**, **T13 Temperate hardwood riparian forest**, **T15 Broadleaved swamp forest on non-acid peat**, **T16 Broadleaved mire forest on acid peat**, **T17 Fagus forest on non-acid soils**, **T18 Fagus forest on acid soils**, **T1B Acidophilous Quercus forest**, **T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T27 Ilex aquifolium forest**, **T31 Temperate mountain Picea forest**, **T32 Temperate mountain Abies forest**, **T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest**, **T35 Temperate continental Pinus sylvestris forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**, **T3F Dark taiga**, **T3G Pinus sylvestris light taiga**, **T3H Larix light taiga**, **T3K Picea mire forest**, **T3M Coniferous plantation of non site-native trees**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz