

Stachys glutinosa

Habitus and growth type

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8**

Moisture indicator value: **3**

Reaction indicator value: **4**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.43**

Disturbance frequency (herb layer): **1.26**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.06**

Grazing pressure: **0.3**

Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S71 Western Mediterranean spiny heath](#)

Constant species of EUNIS habitats: [S71 Western Mediterranean spiny heath](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),**

Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)