Stellaria neglecta

Habitus and growth type

Plant height [m]: **0.36**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **64.1**

Flower

Flowering period: **February-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.73**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.2**
Reaction indicator value: **6.6**
Nutrient indicator value: **6.8**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.44**
Disturbance frequency (herb layer): **1.47**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.47**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.27**
Grazing pressure: **0.19**
Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea*, QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz