# *Stipa dasyvaginata*

### Habitus and growth type

Plant height [m]: **0.6**
Life span: **Perennial**
Life form: **Hemicryptophyte**

### Flower

Flowering period: **May-July**

### Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Disturbance indicator values

Disturbance frequency: **2.24**
Disturbance frequency (herb layer): **2.24**
Disturbance severity: **0.33**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.06**
Grazing pressure: **0.4**
Soil disturbance: **0.12**

### Habitat and sociology

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**

#### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*