Stipa offneri

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Perennial**
Life form: **Hemicryptophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **9.48**

Fruit, seed and dispersal

Seed mass [mg]: **8.1**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7.5**
Moisture indicator value: **3**
Reaction indicator value: **9**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**
Disturbance frequency (herb layer): **1.46**
Disturbance severity: **0.63**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.16**
Grazing pressure: **0.4**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FF (LYG) *Lygeo sparti-Stipetea tenacissimae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**
Constant species of EUNIS habitats: **S61 Western basiphilous garrigue**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz