Stuckenia pectinata

Habitus and growth type

Plant height [m]: **1.43**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **44.91**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **6.17**
Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **6.2**
Moisture indicator value: **11.9**
Reaction indicator value: **7.3**
Nutrient indicator value: **6.6**
Salinity indicator value: **1.7**

Disturbance indicator values

Disturbance frequency: **1.67**
Disturbance frequency (herb layer): **1.67**
Disturbance severity: **0.23**
Disturbance severity (herb layer): **0.23**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.02**
Grazing pressure: **0.01**
Soil disturbance: **0.01**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **NB (POT) *Potamogetonetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **P3a Brackish-water vegetation**, **P3d Fresh-water submerged vegetation**, **P3h Stonewort vegetation**
Constant species of EUNIS habitats: **P3a Brackish-water vegetation**, **P3d Fresh-water submerged vegetation**, **P3h Stonewort vegetation**
Dominant species of EUNIS habitats: **P3d Fresh-water submerged vegetation**

Broad habitat

Occurrence in broad habitats: **Marine, Coastal saltmarsh, Aquatic, Wetland, Saline vegetation**

Distribution

Continentality amplitude: **10**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz