

Stuckenia pectinata

Habitus and growth type

Plant height [m]: **1.43**

Life span: **Perennial**

Life form: **Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **44.91**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **6.17**

Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **6.2**

Moisture indicator value: **11.9**

Reaction indicator value: **7.3**

Nutrient indicator value: **6.6**

Salinity indicator value: **1.7**

Disturbance indicator values

Disturbance frequency: **1.67**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.23**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.02**

Grazing pressure: **0.01**

Soil disturbance: **0.01**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [NB \(POT\) *Potamogetonetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [P3a Brackish-water vegetation](#), [P3d Fresh-water submerged vegetation](#), [P3h Stonewort vegetation](#)

Constant species of EUNIS habitats: [P3a Brackish-water vegetation](#), [P3d Fresh-water submerged vegetation](#), [P3h Stonewort vegetation](#)

Dominant species of EUNIS habitats: [P3d Fresh-water submerged vegetation](#)

Broad habitat

Occurrence in broad habitats: **Marine, Coastal saltmarsh, Aquatic, Wetland, Saline vegetation**

Distribution

Continentality amplitude: **10**