Suaeda acuminata

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.43**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **6.5**
Moisture indicator value: **6.3**

Disturbance indicator values

Disturbance frequency: **2.1**
Disturbance frequency (herb layer): **2.12**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.08**
Grazing pressure: **0.45**
Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***, **ME (THE) *Thero-Salicornietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R64 Semi-desert salt pan**, **S94 Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

riparian scrub

Broad habitat

Occurrence in broad habitats: **Wetland, Saline vegetation, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz