# *Succisella inflexa*

## Habitus and growth type

Plant height [m]: **0.48**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Flower

Flowering period: **July-September**

## Fruit, seed and dispersal

Seed mass [mg]: **1.68**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Temperature indicator value: **6.2**
Reaction indicator value: **5.8**
Nutrient indicator value: **2.8**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.93**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.64**
Grazing pressure: **0.25**
Soil disturbance: **0.12**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**
Continentality amplitude: **1**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*