

# *Tamarix dalmatica*

## Habitus and growth type

Plant height [m]: **3.5**

Life span: **Perennial**

Life form: **Phanerophyte, Tree, Shrub**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **6**

Reaction indicator value: **5**

Nutrient indicator value: **3**

Salinity indicator value: **3**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FD \(NER\) Nerio-Tamaricetea](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Scrub**