

Tanacetum vulgare

Habitus and growth type

Plant height [m]: **0.81**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **20.61**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.21**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Reaction indicator value: **7.1**

Nutrient indicator value: **5.7**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.15**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.7**

Grazing pressure: **0.17**

Soil disturbance: **0.3**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [V38 Dry perennial anthropogenic herbaceous vegetation](#)

Constant species of EUNIS habitats: [V38 Dry perennial anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **8**