

# *Teesdalia coronopifolia*

## Habitus and growth type

Plant height [m]: **0.12**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **January-April**

## Fruit, seed and dispersal

Seed mass [mg]: **0.26**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.4**

Moisture indicator value: **3.4**

Reaction indicator value: **5**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.32**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.44**

Grazing pressure: **0.32**

Soil disturbance: **0.22**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#), [S34 Balkan-Anatolian submontane genistoid scrub](#)

Constant species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#), [S34 Balkan-Anatolian submontane genistoid scrub](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**