

Tephroseris integrifolia

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.56**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **4.4**

Nutrient indicator value: **3.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.14**

Disturbance frequency (herb layer): **1.69**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.29**

Grazing pressure: **0.27**

Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H Larix light taiga](#)

Constant species of EUNIS habitats: [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **7**

Continentality amplitude: **6**