

Teucrium botrys

Habitus and growth type

Plant height [m]: **0.21**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **39.6**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **1.11**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Moisture indicator value: **2.5**

Reaction indicator value: **8**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.24**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.27**

Grazing pressure: **0.3**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**

Continentality amplitude: **2**