

Thesium bergeri

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.74**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.14**

Grazing pressure: **0.35**

Soil disturbance: **0.29**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

Broad habitat

Occurrence in broad habitats: **Scrub**