

# *Thymelaea ruizii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Dwarf shrub**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **2**

Reaction indicator value: **6**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.48**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.14**

Grazing pressure: **0.35**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FP \(ONO\) Festuco hystricis-Ononidetea striatae](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**