

Tofieldia pusilla

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **16.17**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2**

Moisture indicator value: **8.2**

Reaction indicator value: **6.4**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.15**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.25**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.17**

Grazing pressure: **0.14**

Soil disturbance: **0.04**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PA \(SCH\) *Scheuchzerio palustris-Caricetea fuscae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [U25 Boreal and Arctic base-rich scree and block field](#)

Constant species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [U25 Boreal and Arctic base-rich scree and block field](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **6**