

Tordylium maximum

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **24.82**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.38**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **7.6**

Moisture indicator value: **3**

Reaction indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.98**

Disturbance frequency (herb layer): **1.78**

Disturbance severity: **0.73**

Disturbance severity (herb layer): **0.66**

Mowing frequency: **0.61**

Grazing pressure: **0.2**

Soil disturbance: **0.44**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) Sisymbrietea](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality amplitude: **5**