

# *Torilis africana*

## Habitus and growth type

Plant height [m]: **0.57**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.91**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **8**

Moisture indicator value: **4.5**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.54**

Disturbance frequency (herb layer): **1.64**

Disturbance severity: **0.76**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.33**

Grazing pressure: **0.24**

Soil disturbance: **0.31**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**