

# *Tragopogon dubius*

## Habitus and growth type

Plant height [m]: **0.47**

Life span: **Annual, Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **25.22**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **9.2**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **3.6**

Reaction indicator value: **7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **1.82**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.55**

Mowing frequency: **0.61**

Grazing pressure: **0.27**

Soil disturbance: **0.27**

## **Habitat and sociology**

### **EUNIS habitat**

Diagnostic species of EUNIS habitats: [\*\*S34 Balkan-Anatolian submontane genistoid scrub\*\*](#)

Constant species of EUNIS habitats: [\*\*S34 Balkan-Anatolian submontane genistoid scrub\*\*](#)

### **Broad habitat**

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality: **7**

Continentality amplitude: **7**