

Tragopogon ucrainicus

Habitus and growth type

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **5**

Moisture indicator value: **5.8**

Disturbance indicator values

Disturbance frequency: **2.06**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **1.33**

Grazing pressure: **0.27**

Soil disturbance: **0.35**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) Koelerio-Corynephoretea canescentis](#), [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R11 Pannonian and Pontic sandy steppe](#)

Constant species of EUNIS habitats: [S68 Semi-desert sand dune with sparse scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**