

# *Trachynia distachya*

## Habitus and growth type

Plant height [m]: **0.18**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **22.14**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **2.19**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **1.5**

Reaction indicator value: **7**

Nutrient indicator value: **2.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **0.82**

Disturbance frequency (herb layer): **1.65**  
Disturbance severity: **0.71**  
Disturbance severity (herb layer): **0.57**  
Mowing frequency: **0.23**  
Grazing pressure: **0.33**  
Soil disturbance: **0.27**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**