Trifolium dalmaticum

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **7.4**
Moisture indicator value: **2.8**
Reaction indicator value: **6.5**
Nutrient indicator value: **2.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.34**
Disturbance frequency (herb layer): **2.08**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.51**
Grazing pressure: **0.36**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea***, **FJ (TRA) *Stipo-Trachynietea distachyae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz